International Journal of Gastrointestinal Intervention

journal homepage: www.ijgii.org

Case Report

An extremely rare coexistence of achalasia cardia with esophageal leiomyoma: An endoscopic treatment approach

Pankaj Desai^{1,*}, Chintan Patel¹, Mayank Kabrawala², Subhash Nandwani³, Rajiv Mehta³, Ritesh Prajapati³, Nisharg Patel³, and Krishna Parekh⁴

ABSTRACT

The coexistence of an esophageal leiomyoma with an achalasia cardia is extremely rare and poses a diagnostic dilemma as the clinical presentation of an esophageal leiomyoma strongly resembles to an achalasia cardia. Moreover, at most of the time, an esophageal leiomyoma exist with absence of leiomyoma specific symptoms or findings. It could be the reason why the diagnosis of esophageal leiomyoma is missed while the patient has the coexistence of the both. Here, we present a case series of an unusual coexistence of an esophageal leiomyoma with achalasia in three patients who were presented with dysphagia and had an initial diagnosis of achalasia. Even endoscopic evaluation couldn't reveal esophageal leiomyoma; it became apparent during the peroral endoscopic myotomy.

Copyright © 2020, Society of Gastrointestinal Intervention.

Keywords: Achalasia; Leiomyoma; Peroral endoscopic myotomy procedure; Submucosal tunneling endoscopic resection

Introduction

Achalasia is a rare but well-defined neurodegenerative disorder. It is characterised by a defective peristalsis of esophageal body and impaired relaxation of lower esophageal sphincter in response to swallow. It can occur at any age without any gender dominance with the prevalence of 1/100,000 population.¹ Achalasia rarely coexists with other disease conditions. Reported coexisting conditions are pancreatic pseudocyst, mesotheliomas of the pleura, para-esophageal hernia, hiatal hernia, esophageal leiomyoma, and several cancers, such as gastric, pancreatic, hepatic, and esophageal.

The coexistence of achalasia with esophageal leiomyoma is extremely rare due to the insidious onset of leiomyoma.² To the best of our knowledge, only a single case of coexistence is reported till date which was managed laparoscopically.² Herein, we share our experience of an incidental diagnosis of esophageal leiomyoma during the endoscopic management of type I achalasia. The patient was successfully treated using endoscopic approach for leiomyoma.

Case Report

We had three cases of esophageal achalasia coexisting with esophageal leiomyoma. The patients were presented with complaints of dysphagia to solid and liquid. Table 1 summarizes clinical characteristics of the patients. Upper gastrointestinal endoscopy showed fluid, froth, and food residue in the esophageal lumen and tight gastroesophageal junction. Esophageal manometry revealed hypertensive lower esophageal sphincter (LES) and absence of normal peristalsis of esophageal body (Fig. 1). All these findings were suggestive of diagnosis of an achalasia. Hence, peroral endoscopic myotomy (POEM) was scheduled. Patients were intubated and anesthetized. As per the procedure of POEM, esophageal mucosa was lifted away from submucosa by injecting diluted methylene blue dye at 10 to 15 cm proximal to the gastroesophageal junction. An initial longitudinal incision (2–3 cm) was made by triangular tip knife which served as an entry point for the tunnel. The submucosal tunnel was further enlarged by separating mucosa from muscularis using insulated tip knife/hybrid knife. The submucosal tunnel was extended towards the esophago-

¹Department of Endoscopy and Endosonography, Surat Institute of Digestive Sciences, Surat, India

²Department of Gastroenterology, Surat Institute of Digestive Sciences, Surat, India

³Department of Clinical Research, Surat Institute of Digestive Sciences, Surat, India

⁴Department of Gastroenterology, SIDS Hospital and Research Centre, Surat, India

Received May 13, 2020; Revised May 23, 2020; Accepted May 24, 2020

* Corresponding author. Department of Endoscopy and Endosonography, Surat Institute of Digestive Sciences, Majura Gate, Ring Road, Surat 395002, India.

E-mail address: drp_desai@hotmail.com (P. Desai).

Table 1 Clinical Features, Endoscopic Findings, Histopathology, Treatment and Follow-Up of 3 Cases of Esophageal Leiomyoma Coexisted with Achalasia

Variable	Patient no. 1	Patient no. 2	Patient no. 3
Age (yr)/Sex	40/Male	55/Female	36/Female
Clinical features	Dysphagia	Dysphagia	Dysphagia, vomiting
Manometric findings	IRP: 20 mmHg; type I achalasia	IRP: 21.6 mmHg; type I achalasia	IRP: 31.5 mmHg; type I achalasia
Endoscopic findings	Achalasia Cardia	Achalasia Cardia	Achalasia Cardia
Treatment	POEM + STER	POEM + STER	POEM + STER
Histopathologic findings of leiomyoma			
Location	Lower esophagus	Gastroesophageal junction	Gastroesophageal junction
Size (mm)	8	10	6
Histology	Leiomyoma	Leiomyoma	Leiomyoma
Follow-up (3, 6, 12 months)	Asymptomatic	Asymptomatic	Asymptomatic

IRP, integrated relaxation pressure; POEM, peroral endoscopic myotomy; STER, submucosal tunneling endoscopic resection.

Fig. 1. Esophageal manometric examination suggestive of classic pattern of type I achalasia.

gogastric junction; During the dissection, whitish soft tissue lesion was revealed (Fig. 2). This raised the suspicion of presence of leiomyoma. Hence, it was decided to resect the lesion from its surrounding muscular layer using insulated tip knife. Resected tumor was then retrieved out of tunnel using snare through the submucosal entry. The scope was withdrawn from submucosal tunnel. To close the mucosal incision several hemo clips were applied. In view of incidental finding of esophageal leiomyoma, POEM was accompanied with submucosal tunneling endoscopic resection (STER). Histopathological examination (HPE) was suggestive of leiomyoma with immunohistochemistry (IHC) positive for smooth muscle actin, desmin, caldesmon and negative for cluster of differentiation 117. Postoperatively, patients were asymptomatic. On the follow-up of 6, 9, and 12 months, patients were asymptomatic with progressive weight gain.

Discussion

Achalasia is a primary esophageal motility disorder. This condition can be diagnosed by endoscopic and radiological studies, but the main diagnostic modality is manometry. The manometric findings of achalasia are characterized by three features:

an increased integrated relaxation pressure of LES, incomplete relaxation of LES, and aperistalsis of esophagus.¹ Literature search revealed that the coexistence of achalasia with esophageal leiomyoma is extremely rare. The exact incidences are not known but only a single case report is reported in the literature.²

Leiomyoma of esophagus is a benign tumor of the smooth muscle layers. The clinical presentation of esophageal leiomyoma can mimic the symptoms of an achalasia cardia when it is large in size (> 5 cm). Interestingly, an esophageal leiomyoma remains asymptomatic when it is small in size.³ When an esophageal leiomyoma coexist with achalasia, the diagnosis becomes challenging. This is because of the absence of leiomyoma specific symptoms or findings, and the clinical, radiological, and endoscopic findings favoring achalasia cardia. In our case series, clinical picture, endoscopic findings, and esophageal manometry confirmed achalasia. Manometric findings revealed classic pattern of type I achalasia. Even an endoscopic examination of esophagus did not reveal any bulge or mass, probably due to small size of leiomyoma. Thus, in our case series the coexistence of esophageal leiomyoma remained undiagnosed till POEM was performed which highlights the challenges that can be encountered in diagnosis when leiomyoma coexist along with achalasia cardia. Literature review showed several studies which reported a misdiagnosis of achalasia cardia instead of an esophageal leiomyoma mediated dysphagia.^{4,5} Hence, when there is a suspicious bulge or mass on endoscopic findings, an alternative differential diagnosis should be kept in mind, even though the manometric findings are highly suggestive of achalasia cardia. In such cases, we recommend the supplementation of endoscopic ultrasonography (EUS) to aid in further confirmation of final diagnosis. EUS allows visualization of the five layered structure of esophageal wall which may be an added advantage for characterization of the lesion, estimation of size and layer of origin, thus deciding the plan of further management.³

POEM is a novel and minimal invasive endoscopic surgical treatment, widely used in treatment of an achalasia cardia. Inoue et al⁶ was first to demonstrate safety and efficacy of POEM in human for achalasia in 2008. As an extension of POEM technique, STER was developed by Zhou et al.⁷ It is also safe, effective and feasible for treating submucosal tumors originating from submucosa or muscularis propria. Additionally, it provides direct access to the submucosal layer which enables the ease of resection and specimen retrieval for HPE and IHC study.⁷ POEM and STER are two main techniques that are widely used in third space endosco-

Fig. 2. Diagnosis of Esophageal leiomyoma. (A) Histopathology of leiomyoma. (B) Appearance of exposed tumor in the submucosal tunnel during peroral endoscopic myotomy.

py. In the present case series, POEM was planned for treatment of achalasia, but we were fortunate enough to diagnose coexistence of esophageal leiomyoma during POEM. To resect an esophageal leiomyoma, STER was performed with POEM. Thus, the second surgery was avoided.

In conclusion, achalasia cardia is usually idiopathic in origin. Literature search did not show any studies on the relationship between achalasia cardia and leiomyoma. Few case reports showed diffuse leiomyomatosis presented as achalasia cardia. In our case series, leiomyoma was an incidental finding. Further research is warranted to establish a relationship between achalasia and leiomyoma. The coexistence of an achalasia cardia with leiomyoma is a rarely encountered condition and challenging to diagnose pre operatively. POEM and STER remains a valid and effective treatment options.

Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

ORCID

Pankaj Desai, <https://orcid.org/0000-0003-3931-5678>

Chintan Patel, <https://orcid.org/0000-0003-2519-9493>
 Mayank Kabrawala, <https://orcid.org/0000-0003-4707-9695>
 Subhash Nandwani, <https://orcid.org/0000-0001-5648-6837>
 Rajiv Mehta, <https://orcid.org/0000-0003-3197-4337>
 Ritesh Prajapati, <https://orcid.org/0000-0002-8458-276X>
 Nisharg Patel, <https://orcid.org/0000-0002-1183-5826>
 Krishna Parekh, <https://orcid.org/0000-0003-2362-1785>

References

1. Pandolfino JE, Gawron AJ. Achalasia: a systematic review. *JAMA*. 2015;313:1841-52.
2. Palanivelu C, Rangarajan M, Senthilkumar R, Madankumar MV. Laparoscopic surgery for an unusual case of dysphagia: lower oesophageal leiomyoma co-existing with achalasia cardia. *Singapore Med J*. 2008;49:e22-5.
3. Ha C, Regan J, Cetindag IB, Ali A, Mellinger JD. Benign esophageal tumors. *Surg Clin North Am*. 2015;95:491-514.
4. Mohamed A. Education and imaging. Gastrointestinal: pseudoachalasia caused by a lower esophageal stromal tumor. *J Gastroenterol Hepatol*. 2009;24:1152.
5. Ray S, Saluja SS, Gupta R, Chattopadhyay TK. Esophageal leiomyomatosis: an unusual cause of pseudoachalasia. *Can J Gastroenterol*. 2008;22:187-9.
6. Inoue H, Santi EG, Onimaru M, Kudo SE. Submucosal endoscopy: from ESD to POEM and beyond. *Gastrointest Endosc Clin N Am*. 2014;24:257-64.
7. Zhou DJ, Dai ZB, Wells MM, Yu DL, Zhang J, Zhang L. Submucosal tunneling and endoscopic resection of submucosal tumors at the esophagogastric junction. *World J Gastroenterol*. 2015;21:578-83.